

NUB-TH-3147/96
CTP-TAMU-52/96

SUPERSYMMETRIC DARK MATTER

PRAN NATH

*Department of Physics, Northeastern University, Boston
MA 02115, USA*

R. ARNOWITT

*Center for Theoretical Physics, Department of Physics, Texas A & M University,
College Station
TX77843, USA*

A review of recent developments in the analyses of supersymmetric dark matter is given.

1 Introduction

Currently there is a great deal of evidence for the existence of dark matter in the universe: in our galaxy, in other galaxies and in galactic clusters, as much as 90% of matter in the universe may be dark^{1,2,3,4}. There are many candidates for dark matter in astronomy such as jupiters, brown dwarfs, blackholes, etc. However, not all of dark matter can be baryonic and thus a number of particle physics candidates have been suggested. These include massive neutrinos, axions, supersymmetry candidates as well as exotic candidates such as topological defects. A convenient parametrization of matter in the universe is given by the quantity, $\Omega = \rho/\rho_c$, where ρ is the matter density of the universe and $\rho_c = 3 H^2/8\pi G_N$ is the critical matter needed to close the universe, and H is the Hubble parameter which is generally parametrized by $H=100 h$ km/s Mpc. Experimentally the current evaluations of h lie in the interval $0.4 \lesssim h \lesssim 0.8$. An attractive hypothesis theoretically is that of an inflationary universe with $\Omega = 1$. The quantity that enters in theoretical analyses of dark matter is Ωh^2 . The COBE data indicates that there are more than one component to the non-baryonic dark matter, e.g., a hot component comprised of particles which would be relativistic at the time of galaxy formation and a cold component comprised of particles which would be non-relativistic at the time of galaxy formation. The hot component could be a massive neutrino and the cold component could be either axions, or a susy candidate. Actually, in addition to the cold-hot dark matter fit to the power spectrum data observed by COBE and other experiments there are other possibilities as well, such as

inclusion of a cosmological constant. We list below some examples: (i) CHDM (cold-hot DM with $\Omega_{\tilde{\chi}_1^0} = 0.7$, $\Omega_{HDM} = 0.25$, $\Omega_B = 0.05$ (baryonic DM), $h = 0.5$) giving $\Omega_{\tilde{\chi}_1^0} h^2 = 0.18$; (ii) Λ CDM (cosmological constant DM with $\Omega_{\tilde{\chi}_1^0} = 0.25$ – 0.35 , $\Omega_B = 0.05$, $\Omega_\Lambda = 0.6$ – 0.7 , $h = 0.8$) with $\Omega_{\tilde{\chi}_1^0} h^2 = 0.16$ – 0.23 .; (iii) tCMD (tilted DM with $\Omega_{\tilde{\chi}_1^0} = 0.95$, $\Omega_B = 0.05$, $h = 0.5$ – 0.6 , $n = 0.8$) giving $\Omega_{\tilde{\chi}_1^0} h^2 = 0.24$ – 0.34 . In the analysis of this review we shall assume that the parameter space is limited by

$$0.1 \leq \Omega_{\tilde{\chi}_1^0} h^2 \leq 0.4 \quad (1)$$

We will discuss later on the effect of varying the allowed range of $\Omega_{\tilde{\chi}_1^0} h^2$.

Theoretically the analysis of $\Omega_{\tilde{\chi}_1^0} h^2$ is interesting as it involves several different fields of physics: particle physics enters via the fundamental interactions of the dark matter candidates, cosmology enters as the relic density involves both the gravitational constant G_N and the Hubble parameter H , and statistical physics enters via the appearance of the Boltzmann constant k in the analyses of relic density in the early universe. Further, nuclear physics enters as one considers specific detection techniques such as scattering of dark matter particles from nuclei.

In this review our focus will be to explore in detail the supersymmetric candidates for cold dark matter. In the Minimal Supersymmetric Standard Model (MSSM) there are two candidates which are neutral and have odd R parity: these are the neutralino and the sneutrino. If we assume that one of these is the lowest mass supersymmetric particle (LSP) and further assume that R parity is conserved then, one of the candidates could be a CDM candidate. Unfortunately there are 110 arbitrary parameters in MSSM, and thus the masses of the supersymmetric particles are all arbitrary. Thus computations of supersymmetric dark matter in the framework of MSSM are by their very nature highly unpredictable. In contrast, in the minimal supergravity grand unification^{5,6} one can describe low energy physics in terms of just four parameters. Thus the theory is very predictive.

A topic of great interest is the laboratory detection of dark matter. In this context the scattering of dark matter particles from nuclei appears very promising and we will devote considerable amount of discussion to this topic in this paper. The outline of this paper is as follows: In Sec 2 we discuss dark matter in supergravity unification; in Sec 3 a discussion of the relic density analysis is given; in Sec 4 we discuss constraints on dark matter analyses; Sec 5 is devoted to a discussion of the detection of dark matter; in Sec 6 we give an analysis of event rates for the case of universal soft SUSY breaking parameters;

in Sec 7 we discuss the effects of non-universal breaking; in Sec 8 an analysis of event rates with non-universal soft SUSY breaking is given and in Sec 9 we give conclusions.

2 Dark Matter in Supergravity Grand Unification

As discussed above the MSSM is very unproductive. The reason for this is that there is no phenomenologically acceptable mechanism for the breaking of supersymmetry in the MSSM. Consequently one must add by hand the soft SUSY breaking parameters and there are as many as 110 arbitrary parameters that one can add in the soft SUSY breaking sector. For this reason we shall work here within the framework of supergravity grand unification which has a much more limited parameter space. The models we have in mind are those where supersymmetry is broken in the hidden sector and communicated to the physical sector by gravitational interactions, and we consider here the minimal model. We assume that the basic theory in the physical sector is a grand unified theory with a GUT group G which breaks at a scale $M_{GUT} \simeq 10^{16}$ to $SU(3) \times SU(2) \times U(1)$. After the breaking of supersymmetry and the GUT symmetry, and after integration over the fields of the hidden sector and the superheavy fields, one finds the effective potential in the SUSY breaking sector to be of the form:

$$V_{SB} = m_o^2 \Sigma_a z_a^+ z_a + (A_o W_Y + B_o \mu_o H_1 H_2 + h.c) \quad (2)$$

$$L_{mass}^\lambda = -m_{1/2} \bar{\lambda}^\alpha \lambda^\alpha \quad (3)$$

where $\{z_a\}$ are the scalar fields, λ^α the gaugino fields, μ is the Higgs mixing parameter and W_Y is the cubic Yukawa coupling which enters in the effective superpotential, W , of the low energy theory as follows:

$$W = \mu_o H_1 H_2 + W_Y + \frac{1}{M_G} W^{(4)} \quad (4)$$

Here $W^{(4)}$ contains any quartic non-renormalizable couplings which may lead to proton decay. At this level the theory contains the four SUSY breaking parameters m_o , $m_{1/2}$, A_o , and B_o in addition to the Higgs mixing parameter μ_o , and the GUT parameters M_G and α_G .

The dynamics of supergravity grand unification breaks the electro-weak symmetry dynamically by radiative effects^{7,8,9}. Further, one typically uses the radiative breaking of the electro-weak symmetry to compute the Higgs mixing parameter μ by fixing the Z mass, and one eliminates the parameter B (which is the value of the parameter B_o at the electro-weak scale) in favor of $\tan\beta \equiv \langle H_2 \rangle / \langle H_1 \rangle$ such that

$$\mu^2 = \frac{\mu_1^2 - \mu_2^2 \tan^2 \beta}{\tan^2 \beta - 1} - \frac{1}{2} M_Z^2; \quad \sin 2\beta = \frac{-2B\mu}{2\mu^2 + \mu_1^2 + \mu_2^2} \quad (5)$$

Here $\mu_i^2 = m_{H_i}^2 + \Sigma_i$ where $m_{H_i}^2$ is the running H_i mass at scale $Q \approx M_Z$ and Σ_i are loop corrections. The $m_{H_i}^2$ are given by

$$m_{H_1}^2 = m_o^2 + m_{1/2}^2 g(t) \quad (6)$$

$$m_{H_2}^2 = m_{1/2}^2 e(t) + A_o m_{1/2} f(t) + m_o^2 h(t) - k(t) A_o^2 \quad (7)$$

where the form factors e,f,g,h,k are defined in ref(10) and $t = \ln(M_G^2/Q^2)$. One may show that solutions exist to Eqs. (5), i.e. that $SU(2) \times U(1)$ is spontaneously broken, if and only if at least one of the supersymmetry soft breaking interactions are non-zero. Thus it is the supergravity interactions at M_G that give rise to the breaking of $SU(2) \times U(1)$ at the electroweak scale. The $SU(3) \times SU(2) \times U(1)$ gaugino masses at the electro-weak scale are given by

$$m_i = (\alpha_i/\alpha_G) m_{1/2} \quad (8)$$

In the above supergravity grand unification the sparticle mass spectrum is determined dynamically in terms of the four parameters, and thus the theory predicts what the LSP is. It is quite remarkable that over most of the parameter space one finds that the LSP is indeed a neutralino $\tilde{\chi}_1^0$. Thus supergravity predicts the $\tilde{\chi}_1^0$ to be the cold dark matter of the universe. The $\tilde{\chi}_1$ is of course an admixture of gauginos $\tilde{W}_3, \tilde{B}$ and Higgsinos $\tilde{H}_1, \tilde{H}_2$ and one may write

$$\tilde{\chi}_1 = n_1 \tilde{W}_3 + n_2 \tilde{B} + n_3 \tilde{H}_1 + n_4 \tilde{H}_2 \quad (9)$$

where we use the convention that $\tilde{\chi}_1$ is the lightest neutralino and where n_i are the components of the gaugino states and are to be determined by diagonalizing the neutralino mass matrix. The neutralino mass matrix in the basis $(\tilde{W}_3, \tilde{B}, \tilde{H}_1, \tilde{H}_2)$ is given as follows:

$$M_{\tilde{Z}} = \left(\begin{array}{cc|cc} \tilde{m}_2 & o & a & b \\ o & \tilde{m}_1 & c & d \\ \hline a & c & o & -\mu \\ b & d & -\mu & o \end{array} \right) \quad (10)$$

Here $a = M_Z \cos \theta_W \cos \beta$, $b = -M_Z \cos \theta_W \sin \beta$, $c = -M_Z \sin \theta_W \cos \beta$ and $d = M_Z \sin \theta_W \sin \beta$. In general one needs a numerical analysis to determine the value of the mass eigenvalues as well as the eigenvectors. In the limit $\mu^2 \gg M_Z^2$, which holds in a significant part of the parameter space, it is possible to obtain an analytic form for the indices n_i as an expansion in μ/M_Z .

The results are given in appendix A. This expansion is useful in understanding the phenomena associated with the neutralino relic densities and event rates. Generally one finds that in this limit the neutralino is mostly a Bino with $n_2 > 0.95$ and often larger, while n_3, n_4 and n_1 are typically of first order, $O(M_Z/\mu)$. Because of this the Higgsino components are not totally negligible, and they are often quite significant e.g. $n_3 \approx 0.2$. Hence one cannot neglect the Higgsino components even when the neutralino may be dominantly a Bino. This result has important implications in the analysis of event rates.

Before proceeding further we wish to discuss the effect that a heavy top quark has on the electro-weak breaking. One of the major effects of a heavy top arises because it lies close to the Landau pole value which is given by $m_t^f \approx 197 \sin \beta$. This occurs because the proximity of the top to the Landau pole drives the lighter stop towards a tachyonic limit. We exhibit this phenomenon in some detail. The mass of the lighter stop $\tilde{t}_1$ is determined by the diagonalization of the following stop $(mass)^2$ matrix:

$$\begin{pmatrix} m_{\tilde{t}_R}^2 & m_t(A_t - \mu c t n \beta) \\ m_t(A_t - \mu c t n \beta) & m_{\tilde{t}_L}^2 \end{pmatrix} \quad (11)$$

The lighter stop $(mass)^2$ eigen-value in the vicinity of the Landau pole is given by¹¹

$$m_{\tilde{t}_1}^2 = -\frac{1}{3} \frac{A_R^2}{D_o} + m_{\tilde{t}_1}^2(NP) \quad (12)$$

where

$$A_R \cong A_t - 0.613 m_{\tilde{g}}; \quad D_o \cong 0.164 [(m_t^f)^2 - (m_t)^2] / M_W^2 \quad (13)$$

and where $m_{\tilde{t}_1}^2(NP)$ is a non-pole contribution. The condition that $\tilde{t}_1$ not turn tachyonic limits severely the allowed range of A_t .

3 Relic Density Analysis

An important quantity in the analysis of supersymmetric dark matter is the LSP relic density. After the time of the Big Bang, the primordial neutralinos begin to annihilate into ordinary matter, e.g., via the process $\tilde{\chi}_1^0 + \tilde{\chi}_1^0 \rightarrow f + \bar{f}, \dots$, where $f, \bar{f}$ are leptons, quarks, etc¹². The annihilation is governed by the Boltzman equation :

$$\frac{dn}{dt} = -3Hn - \langle \sigma v \rangle (n^2 - n_0^2) \quad (14)$$

where n = number/vol is the $\tilde{\chi}_1^0$ number density, n_0 is the value of n at thermal equilibrium, H is the Hubble parameter at time t , σ is the neutralino annihilation cross section, v is the relative neutralino velocity, and $\langle \sigma v \rangle$ stands for the thermal average. Assuming that the neutralinos are non-relativistic at the time of interest (a realistic approximation) one has

$$\langle \sigma v \rangle = \int_0^\infty dv v^2 \sigma v \exp[-v^2/4x] / \int_0^\infty dv v^2 \exp[-v^2/4x] \quad (15)$$

At large temperatures close to the Big Bang the $\tilde{\chi}_1^0$ is in thermal equilibrium with the background. However, the $\tilde{\chi}_1^0$ decouples from the background when the annihilation rate becomes smaller than the expansion rate given by the Hubble term. This decoupling takes place at the "freezeout" temperature T_f . Integration of the Boltzman equation from the freezeout to the current temperatures gives the current relic density. One finds

$$\Omega_{\tilde{\chi}_1^0} h^2 \cong 2.48 \times 10^{-11} \left(\frac{T_{\tilde{\chi}_1^0}}{T_\gamma} \right)^3 \left(\frac{T_\gamma}{2.73} \right)^3 \frac{N_f^{1/2}}{J(x_f)} \quad (16)$$

where

$$J(x_f) = \int_0^{x_f} dx \langle \sigma v \rangle(x) \text{GeV}^{-2} \quad (17)$$

and $x_f = kT_f/m_{\tilde{\chi}_1}$. Now numerical analyses show that freezeout occurs for values of $x_f \sim \frac{1}{20}$ which indeed corresponds to the neutralinos being non-relativistic at the time of the freezeout. The central quantity in the analysis of the relic density is $J(x_f)$. The computation of this quantity is quite subtle as it involves a double integral over (σv) which in some regions is rapidly varying. An approximation that has generally been made is to expand σv in a power expansion in v , i.e. $\sigma v \cong a + b(v^2/c^2) + \dots$. With this approximation the computation of $J(x_f)$ becomes trivial. However, it is known¹³ that such an expansion is not a good approximation when (σv) is rapidly varying. This is precisely the situation that occurs in the annihilation of the neutralinos when one is close to the Z and the Higgs poles^{14,15,16}. In this case one must carry out a thermal averaging keeping the pole structure intact. To circumvent the problem of the double integration over the pole, one can carry out analytically one of the integrations reducing $J(x_f)$ to a single integration which is computationally easy to evaluate. Details are given in Appendix B.

4 Constraints on Dark Matter Analyses

The analysis of dark matter relic density is subject to many constraints. Since the analysis is carried out in the framework of supergravity grand unification one computes masses of 32 supersymmetric particles in terms of the basic SUSY breaking parameters (which are just four for the universal soft SUSY breaking case). The analysis of the mass spectrum is subject to following constraints: (i) color and charge conservation at the electro-weak scale, (ii) naturalness constraints, i.e., that $m_0, m_{\tilde{g}} \leq 1\text{TeV}$, (iii) LEP, CDF and DO limits on the mass spectrum. Further, the decay $b \rightarrow s + \gamma$ represents an important constraint on any new physics beyond the Standard Model. Recently CLEO measured the branching ratio for $B \rightarrow X_s + \gamma$ to be⁷:

$$BR(B \rightarrow X_s \gamma) = (2.32 \pm 0.5 \pm 0.29 \pm 0.32) \times 10^{-4} \quad (18)$$

Combining the errors in quadrature gives $BR(B \rightarrow X_s \gamma) \cong (2.32 \pm 0.66) \times 10^{-4}$. In the spectator approximation one can relate the above to the flavor changing process $b \rightarrow s + \gamma$ and one has

$$\frac{BR(B \rightarrow X_s \gamma)}{BR(B \rightarrow X_c e \bar{\nu}_e)} \cong \frac{\Gamma(b \rightarrow s + \gamma)}{\Gamma(b \rightarrow c + e + \bar{\nu}_e)} \equiv R \quad (19)$$

where $BR(B \rightarrow X_c e \bar{\nu}_e) = (10.7 \pm 0.5)$. The quantity R turns out to be a useful parameter as the factor (m_b^5) (which contains a significant experimental ambiguity) as well as some CKM factors cancel out in this ratio.

The $b \rightarrow s + \gamma$ decay proceeds via loop corrections. For the case of the Standard Model the loop contribution involves a W - t exchange. For the supersymmetric case one has in addition a $\tilde{W} - \tilde{t}$, $H^- - t$ and a $\tilde{Z} - \tilde{b}$ exchange. We see then that the SUSY contributions enter on an equal footing with the Standard Model contribution. Thus $b \rightarrow s + \gamma$ is an important constraint on the SUSY parameter space. The supersymmetric contribution to the decay $b \rightarrow s + \gamma$ is governed by effective Hamiltonian¹⁸

$$H_{eff} = V_{tb} V_{ts}^* \frac{G_F}{\sqrt{2}} C_7(M_W) Q_7 \quad (20)$$

where $F_{\mu\nu}$ is the electromagnetic field strength and $Q_7 = (e/24\pi^2) m_b \bar{s}_L \sigma^{\mu\nu} b_R F_{\mu\nu}$. However, the computation of $b \rightarrow s + \gamma$ must be carried out at the scale $\mu \simeq m_b$ where the decay occurs and thus one must use the renormalization group equations to go from the scale $\mu \approx M_W$ to $\mu \simeq m_b$. Using the above evolution, the value of R to leading order (LO) in QCD corrections is given by

$$R = \left| \frac{V_{tb} V_{ts}^*}{V_{cb}} \right|^2 \frac{6\alpha}{\pi I(z)} |C_7^{eff}(m_b)|^2 \quad (21)$$

where $I(z) = 1 - 8z^2 + 8z^6 - z^8 - 24z^4 \ln z$ is a phase space factor ($z = m_c/m_b$) and $C_7^{eff}(m_b)$ is given by

$$C_7^{eff}(m_b) = C_2(M_W) + \eta^{\frac{16}{23}} C_7(M_W) + \frac{8}{3} (\eta^{\frac{14}{23}} - \eta^{\frac{16}{23}}) C_8(M_W) \quad (22)$$

Here $C_8(M_W)$ is the co-efficient of the operator $Q_8 = (g_3/16\pi^2) m_b \bar{s}_R \sigma^{\mu\nu} T^A b_L G_{\mu\nu}^A$ in the renormalization group evolution, T^A and $G_{\mu\nu}^A$ are the gluon generators and field strengths, and C_2 represents the operator mixing with the 4-quark operators in the renormalization group evolution.

The Standard Model leading order prediction gives¹⁹ $BR[B \rightarrow X_s \gamma] \cong (2.9 \pm 0.8) \times 10^{-4}$ for $m_t = 174$ GeV. Several analyses of $b \rightarrow s + \gamma$ in SUSY models within the leading order(LO) approximation have been carried out, and their effects on dark matter analysed. Generally one finds that the $b \rightarrow s + \gamma$ constraint is very stringent for $\mu > 0$ but less so for the case $\mu < 0$. Thus for $\mu > 0$ one finds that as much as 2/3 of the parameter space is eliminated while for $\mu < 0$ only about 1/5 of the parameter is eliminated^{20,21,32,33}. More recently, the next to leading order corrections to the Standard Model have been computed²². Analyses including these give results which are similar²³ to the ones based on the LO corrections^{20,21,32,33}.

5 Detection of Dark Matter

A variety of possibilities for the detection of dark matter which involves more than just its gravitational effects have been discussed in the literature. These include :(i) scattering by nuclei of terrestrial detectors ,(ii) annihilation in the centre of sun or earth, (iii) annihilation in the halos of galaxies, (iv) atomic excitations, and (v) superheated microbubbles. Presently, the most promising of the above are (i) and (ii), and we shall discuss specifically (i) in detail here. Case(ii) is based on the idea that neutralinos in the solar neighborhood will be attracted to the center of the sun, where over a period of time they will lose energy and accumulate. Annihilation of these will produce neutrinos which in the terrestrial detection apparatus will show up as upward moving muons(via their neutral current interaction). A similar situation is expected for the annihilation in the center of the earth. Background for these processes are significantly reduced because of the small angular apertures around the center of sun and earth^{24,25}. Signals of this type can be picked up by neutrino telescopes. In case (iii) one expects that neutralinos in the halos of galaxies will annihilate into $(\gamma, e^+, \bar{p}, \dots) + X$. However, the signal to background ratio does not appear very promising in this case. Case(iv) involves scattering of neutralinos by electrons bound in atoms which may result in atomic excitations²⁶.

Realistic analyses of this possibility are currently lacking. Finally we mention the possibility of superheated microbubbles as a cold dark matter detector which has been discussed recently²⁷.

We turn now to a detailed discussion of neutralino-nucleus scattering^{28–38}. The fundamental interaction here is the neutralino-quark scattering. However, since the quarks are bound inside the nucleon and the nucleons are bound inside the nucleus, one has to deal with nuclear physics effects. We shall return to these later. We consider first the fundamental interaction involving neutralino and quarks. This interaction involves exchange of Z boson, squarks and Higgs. The dark matter in the Milky Way halo is estimated to have a velocity of $v \sim 300 \text{ km/s}$. In scattering involving these neutralinos the typical kinetic energy exchange is $O(1 \text{ keV } M_{\chi_1}/1 \text{ GeV})$. Because the energy exchanges are small one can approximate the pole exchange diagrams by contact interactions. The total interaction is of the form

$$L_{eff} = L_{eff}^{(SD)} + L_{eff}^{(SI)} \quad (23)$$

where $L_{eff}^{(SD)}$ is a spin dependent interaction and $L_{eff}^{(SI)}$ is a spin independent part. $L_{eff}^{(SI)}$ is given by

$$L_{eff}^{(SD)} = (\bar{\chi}_1 \gamma^\mu \gamma^5 \chi_1) [\bar{q} \gamma_\mu (A_L P_L + A_R P_R) q] \quad (24)$$

where $P_{R,L}$ are projection operators with $P_{R,L} = (1 \pm \gamma_5)/2$ etc. Note that because of the Majorana nature of the neutralino $\bar{\chi}_1 \gamma^\mu \chi_1 = 0$ and thus only the axial-vector bilinear involving the neutralino enters. The spin-dependent interaction receives contributions from the Z exchange and from the chirality diagonal (LL and RR) squark exchange terms. For the Z-exchange one finds that $A_L^Z, A_R^Z \propto (n_3^2 - n_4^2)$ which means that this exchange not only depends on the Higgsino components of the neutralino [defined in Eq.(9)] but on their mismatch as well. On the other hand the chirality diagonal squark exchange depends on the gaugino components of the neutralino, i.e., one has

$$A_L^{\tilde{q}} \propto (n_1 g_2 T_3 + n_2 g_1 \frac{Y_L}{2})^2, \quad A_R^{\tilde{q}} \propto (n_2 g_1 \frac{Y_R}{2})^2 \quad (25)$$

We note in passing that the spin dependent interaction discussed above involves no interference between the gaugino components and the Higgsino components. This aspect of the spin dependent interaction is in contrast to the scalar interaction which we discuss next. The effective scalar interaction is given by

$$L_{eff}^{(SI)} = (\bar{\chi}_1 \chi_1) (\bar{q} C m_q q) \quad (26)$$

The scalar interaction receives contributions from Higgs exchange and from the squark exchange. The contribution to C from the Higgs exchange obeys

$$C_{Higgs} \propto \frac{g_2^2}{4M_W} \left[\frac{F_h}{m_h^2} \left\{ \frac{\cos\alpha_H}{\sin\beta} \right\} + \frac{F_H}{m_H^2} \left\{ \frac{\sin\alpha_H}{\sin\beta} \right\} \right] \begin{matrix} u - quark \\ d - quark \end{matrix} \quad (27)$$

where h is the light CP even Higgs, H is the heavy CP even Higgs, $F_h = (n_1 - n_2 \tan\theta_W)(n_4 \cos\alpha_H + n_3 \sin\alpha_H)$, $F_H = (n_1 - n_2 \tan\theta_W)(n_4 \sin\alpha_H - n_3 \cos\alpha_H)$ and α_H is the rotation angle needed to diagonalize the h-H⁰ mass matrix and is defined such that³⁹

$$\cos(2\alpha_H) = \frac{M_{11}^H - M_{22}^H}{\sqrt{(M_{11}^H - M_{22}^H)^2 + 4M_{12}^H}}, \quad \tan 2\alpha_H = \frac{2M_{12}^H}{(M_{11}^H - M_{22}^H)} \quad (28)$$

where M_{ij}^H is the Higgs mass matrix in the CP even sector. We emphasize here the fact that the inclusion of loop corrections to the Higgs masses is important in the computation of the Higgs mixing angle^{40,31}. Further, the contribution of the heavy Higgs exchange is in general not small⁴¹. Thus while the heavy Higgs contribution is indeed small in the up quark sector, its contribution in the down quark sector can be very significant. The reason for this is that in the down quark sector the light Higgs contribution is proportional to $\sin\alpha_H$ while the heavy Higgs contribution is proportional to $\cos\alpha_H$. Now over much of the parameter space one finds that $\tan\alpha_H = O(1/10)$. This leads to a suppression in the light Higgs sector and an enhancement in the heavy Higgs sector. Further, $n_3/n_4 \sim \tan\beta$ (see Appendix A) and hence F_H can become significantly larger than F_h . These enhancements in the heavy sector can sometimes even overcome the natural mass suppression of m_h^2/m_H^2 making the heavy Higgs sector contribution quite large. Finally, the (chirality non-diagonal) squark exchange also contributes to the scalar interaction and one has:

$$C_{\bar{q}} \propto \left(\frac{n_1 g_2 T_3 + n_2 g_1 \frac{Y_L}{2}}{(m_{\bar{q}_L}^2 - m_{\chi_1}^2)} - \frac{n_2 g_1 \frac{Y_R}{2}}{(m_{\bar{q}_L}^2 - m_{\chi_1}^2)} \right) \left(n_3 \frac{T_3 + \frac{1}{2}}{\sin\beta} - n_4 \frac{T_3 - \frac{1}{2}}{\cos\beta} \right) \quad (29)$$

We note that the scalar interaction involves an interference of the gaugino and the Higgsino components of the neutralino. Thus if the lightest neutralino were to be strictly a Bino, the scalar interaction would vanish and the neutralino- nucleus scattering would be strictly governed by the spin dependent interaction. However, such a situation is never realised and the scalar

interaction always plays an important role in the event rate analysis. The scalar interaction is further enhanced by the fact that when one sums over all the quarks in the nucleus, one produces in the case of the scalar interaction a nuclear mass factor M_N , which enhances the contribution of the scalar term. Thus the scalar interaction is the dominant one for heavy nuclei. The event rate for a detector involving a target nucleus of mass M_N is then given by

$$R = [R_{SI} + R_{SD}] \left[\frac{\rho_{\tilde{\chi}_1}}{0.3 \text{ GeV cm}^{-3}} \right] \left[\frac{v_{\tilde{\chi}_1}}{320 \text{ km/s}} \right] \frac{\text{events}}{\text{kg da}} \quad (30)$$

Here $\rho_{\tilde{\chi}_1}$ is the local mass density of $\tilde{\chi}_1$, and $v_{\tilde{\chi}_1}$ is the incident $\tilde{\chi}_1$ velocity. The spin dependent (SD) rate is given by

$$R_{SD} = \frac{16m_{\tilde{\chi}_1}M_N}{[M_N + m_{\tilde{Z}_1}]^2} \lambda^2 J(J+1) |A_{SD}|^2 \quad (31)$$

where J is the nuclear spin and λ is defined by $\langle N | \sum \vec{S}_i | N \rangle = \lambda \langle N | \vec{J} | N \rangle$. For the spin independent part (SI) one has

$$R_{SI} = \frac{16m_{\tilde{\chi}_1}M_N^3M_Z^4}{[M_N + m_{\tilde{\chi}_1}]^2} |A_{SI}|^2 \quad (32)$$

We note that for large M_N , $R_{SD} \sim 1/M_N$, while $R_{SI} \sim M_N$ which implies that heavier the nucleus the more sensitive it is to the spin independent scattering.

We discuss now the uncertainties associated with the computation of R_{SI} and R_{SD} . In the computation of R_{SI} most of the uncertainty is associated with the computation of the matrix elements of the operator $m_q \bar{q}q$ between the nucleon states. One may write the matrix elements in the form

$$\langle n | m_q \bar{q}q | n \rangle = m_n f_q^{(n)} \quad (33)$$

Estimates give $f_u \approx f_d \approx 0.05$, $f_s \approx 0.2$ with as such much as 50% uncertainty in their determination⁴². Further the heavy quark contributions cannot be ignored. They couple to nucleons at the loop level via exchange of gluons and contribute to the mass of the nucleon. One may write $\langle n | m_Q \bar{Q}Q | n \rangle = \frac{2}{27} m_n (1 - \sum_q f_q^{(n)})$. An uncertainty also exists in the determination of R_{SD} . This quantity depends on the matrix element

$$\langle n | \bar{q} \gamma^\mu \gamma^5 q | n \rangle = 2s_{(n)}^\mu \Delta q \quad (34)$$




Figure 1: Maximum and minimum of event rates for Ge with scalar interaction.

Here $s_{(n)}^\mu$ is the spin 4-vector of the nucleon, and Δq is the quark polarizability. Currently there are two determinations of Δq , one using the older EMC data⁴³, and the other using the new SMC data⁴⁴. The analyses using the EMC data and the SMC data give

$$\Delta u = 0.7 \pm 0.08, \Delta d = -0.49 \pm 0.08, \Delta s = -0.15 \pm 0.08(EMC) \quad (35)$$

$$\Delta u = 0.83 \pm 0.03, \Delta d = -0.43 \pm 0.03, \Delta s = -1.0 \pm 0.03(SMC) \quad (36)$$

Now the variations in Δq , and specifically the variation in Δs can give rise to a significant variation in R_{SI} . However, R_{SI} is typically not a large component of the total R except for light nuclei such as CaF_2 . Thus the event rate analyses in lighter nuclei are the ones most sensitive to variations in the values of Δq while the event rates in heavier nuclei are not very sensitive.

6 Analysis of Event Rates for Universal Soft SUSY Breaking

We discuss now the analysis of the event rates. Here we give the analysis under the assumption that the soft SUSY breaking parameters are universal at the GUT scale. As discussed in Sec 2, the parameter space of supergravity grand unification in this case is 4 dimensional and is spanned by

$$m_0, m_{\tilde{g}}, A_t, \tan\beta, \text{sign}(\mu) \quad (37)$$

As mentioned already in sec 2 the condition that $\tilde{t}_1$ not turn tachyonic places severe constraints on the parameter space. From eqs(12) and (13) one may see




Figure 2: Maximum and minimum of event rates for Ge with spin dependent interaction.

this as follows: for A_t negative A_R^2 becomes large quickly since the A_t term and the gluino term reinforce each other driving the $\tilde{t}_1$ tachyonic. For A_t positive one finds that A_t also cannot get too large since that would make A_R^2 large ($m_{\tilde{g}} < 1TeV$) and again drive the light stop tachyonic. Numerical analysis shows that the no tachyonic condition limits A_t to the following range^{11,33}

$$-1.0 \lesssim A_t/m_o \lesssim 5.5 \quad (38)$$

Thus a heavy top mass eliminates a large amount of the SUSY parameter space.

Event rates are computed under the constraint of the naturalness assumption on $m_0, m_{\tilde{g}} \leq 1TeV$, and $\tan\beta < 25$. A_t is then restricted by the electro-weak symmetry breaking and by the condition that there be no tachyons, i.e., eq(38). In this range of the parameter space we compute the event rates for various target materials. The maximum and the minimum of the event rates for Ge are exhibited in Figs 1-3. Fig.1 is for coherent scattering when we take into account only the scalar part of the interaction. Fig.2 gives the analysis of incoherent part of the event rates where we take into account only the spin dependent part. Fig.3 gives a composite of the event rates including the coherent part, the incoherent part and the total. A comparison of the maximum and the minimum curves shows that for the case of Ge (a heavy nucleus) the maximum event rates are almost 100% dominated by the coherent part, while the minimum curves have a substantial component (as much as 40%) of incoherent part above a neutralino mass of about 60 GeV. Fig.4 shows the corresponding curves for a light nucleus, CaF_2 . We also note that both the maximum and minimum event rates show a sharp dip in the vicinity of the Z-pole. This phenomenon arises as a consequence of the accurate method of computing the relic density which takes account of the correct thermal aver-

aging over the Z and the Higgs poles in the annihilation of the neutralinos (the lack of a second sharp dip in the event rates from the Higgs pole is due to the fact that one is integrating over the parameter space which smears the Higgs mass pole effects). The dependence of the event rates on the neutralino mass beyond the annihilation region involving the Z and the Higgs poles can also be understood. The event rates in this region are controlled by μ . Since μ^2 is an increasing function of $-m_{H_2}^2$, by Eq. (7) it is an increasing function of $m_{\tilde{\chi}_1}$ and m_o . As discussed above the maximum event rates are dominated by A_{SI} which is proportional to $(n_2 n_3)$ [or $(n_2 n_4)$] which by Eqs. (48-51) are the leading $[O(M_Z/\mu)]$ terms. These decrease as μ increases. Thus we expect R to decrease as $m_{\tilde{\chi}_1}$ and m_o increase. The analysis of Fig 3 bears this out. Finally we mention that the the maximum curve in the event rates has a strong dependence on $\tan\beta$. It increases as $\tan\beta$ increases. This can be seen from Eq.(27) where the leading d-quark contribution has $C_{Higgs} \sim 1/\cos\beta \sim \tan\beta$.

7 Effects of Non-universal Soft Breaking

Most of the previous analyses in supergravity grand unification have been done using universal boundary conditions for the soft SUSY breaking masses at the scale M_G . However, the framework of supergravity unification allows for the presence of non-universalities from a general Kahler potential⁴⁵. In the present analysis we wish to examine the effect of non-universalities on the predictions on dark matter event rates. In the present analysis for specificity we shall confine ourselves to consideration of non-universalities only in the Higgs sector^{46,47,48,49}. Thus we assume that the only non-universalities we have at the GUT scale M_G are given by

$$m_{H_1}^2 = m_0^2(1 + \delta_1); \quad m_{H_2}^2 = m_0^2(1 + \delta_2), \quad (39)$$

where $\delta_{1,2}$ represent the deviation from the universal reference mass m_0 . All the remaining parameters are assumed universal. The important parameters that enter in the low energy domain are μ^2 , m_A^2 , and $m_{\tilde{t}_{1,2}}$. The role of $m_{\tilde{t}_{1,2}}$ has already been discussed in Sec.2 We exhibit here the effects of non-universalities on the parameters μ^2 and m_A^2 . We find

$$\mu^2 = m_0^2 C_1 + A_0^2 C_2 + m_{\frac{1}{2}}^2 C_3 + A_0 m_{\frac{1}{2}} C_4 - \frac{1}{2} M_Z^2 \quad (40)$$

Here

$$C_1 = \frac{1}{t^2 - 1} \left(1 - \frac{3D_0 - 1}{2} t^2\right) + \frac{1}{t^2 - 1} \left(\delta_1 - \frac{D_0 + 1}{2} \delta_2 t^2\right) \quad (41)$$




Figure 3: A comparison of the maximum and the minimum of event rates for Ge for three cases:(i) the coherent part arising from the scalar interaction (dotted),(ii) the incoherent part arising from the spin dependent interaction(dashed), and (iii) sum of cases(i) and (ii) (solid). The maximum dotted line overlaps the maximum solid line(see Fig 1 for comparison of the dotted and solid lines)

$$C_2 = -\frac{t^2}{t^2 - 1}k, \quad C_3 = -\frac{1}{t^2 - 1}(g - t^2e), \quad C_4 = -\frac{t^2}{t^2 - 1}f \quad (42)$$

where $t \equiv \tan\beta$ and where the functions e,f,g,k are as defined in ref(10). Similarly, for m_A^2 one has

$$m_A^2 = m_0^2 D_1 + A_0^2 D_2 + m_{\frac{1}{2}}^2 D_3 + A_0 m_{\frac{1}{2}} D_4 - \frac{1}{2} M_Z^2 \quad (43)$$

where

$$D_1 = \frac{t^2 + 1}{t^2 - 1} \left(1 - \frac{3D_0 - 1}{2}\right) + \frac{t^2 + 1}{t^2 - 1} \left(\delta_1 - \frac{D_0 + 1}{2} \delta_2 t^2\right) \quad (44)$$

$$D_2 = -\frac{t^2 + 1}{t^2 - 1}k, \quad D_3 = -\frac{t^2 + 1}{t^2 - 1}(g - e), \quad D_4 = -\frac{t^2 + 1}{t^2 - 1}f \quad (45)$$

In the above D_0 defines the Landau pole in the top quark Yukawa coupling y_0 , i.e.,

$$y_0 = \frac{y_t}{E(t)D_0}; \quad D_0 = 1 - 6y_t \frac{F(t)}{E(t)} \quad (46)$$




Figure 4: A comparison of the maximum and minimum of event rates for CaF_2 for three cases:(i) the coherent part arising from the scalar interaction (dotted),(ii) the incoherent part arising from the spin dependent interaction(dashed), and (iii) sum of cases(i) and (ii) (solid)

and

$$E(t) = (1 + \beta_3)^{\frac{16}{3b_3}} (1 + \beta_2)^{\frac{3}{b_2}} (1 + \beta_1)^{\frac{13}{9b_1}} \quad (47)$$

where $\beta_i = \alpha_i(0)b_i/4\pi$, $\alpha_i(0)$ are the gauge coupling constants at the GUT scale with $\alpha_1 = (5/3)\alpha_Y$, b_i are the one loop beta function co-efficients defined by $(b_1, b_2, b_3) = (33/5, 1, -3)$, and $F(t) = \int_0^t E(t)dt$.

Normally we expect $|\delta_{1,2}| \leq 0.5$, and a similar level of non-universality in the relevant mass spectra at the electro-weak scale. However, under special circumstances enhancement of non-universalities can occur when the universal terms are suppressed. To exhibit this phenomenon we consider the solution for μ^2 . The dominant terms here are the first two terms, i.e., the terms which involve C_1 and C_2 . Now C_1 contains a universal part and a non-universal part. For large $\tan\beta$, the universal part accidentally cancels when $D_0=1/3$ which occurs for a value of $m_t \approx 168$ GeV, and consequently the non-universality effects get enhanced. An enhancement of non-universalities also occurs from the $A_0 C_2$ term, but in a somewhat different manner. Here the non-universalities are enhanced when the residue of the Landau pole in A_0 vanishes. This happens when $A_t \approx 0.6 m_{\tilde{g}}$.




Figure 5: Maximum and minimum of event rates for Xenon for the universal case when $\delta_1 = 0 = \delta_2$

8 Event Rates With Non-universal Soft SUSY Breaking

In this section we analyse the event rates for Xenon with both universal as well as with non-universal soft SUSY breaking terms. Fig.5 exhibits the maximum and the minimum of event rates for Xenon for the case $\delta_1 = 0 = \delta_2$. The general features of this case are similar to the case of Ge discussed earlier and the observations made there apply here as well. Next we discuss the non-universal case. When $\delta_2 > 0$ and $\delta_1 < 0$, the non-universal terms in Eq.(40) decrease μ^2 and can in some cases drive μ^2 negative, eliminating such points from the parameter space. This effect is larger for small $\tan\beta$. Fig.6 exhibits this case for $\delta_2 = -\delta_1 = 1$ where the major effect is to significantly raise the minimum event rate (for lower values of $m_{\tilde{\chi}_1}$) since it is the smallest $\tan\beta$ that gives rise to the minimum values of R . Fig.7 exhibits the opposite effect, i.e., a decrease in minimum value of R , when $\delta_2 = -1$, $\delta_1 = 1$. The decrease in the minimum arises since μ^2 is increased and R is a falling function of μ . Fig 8 gives for comparison the composite of the three cases discussed above. We note that as $m_{\tilde{\chi}_1}$ increases and hence $m_{\tilde{g}}$ increases one finds that A_R increases and hence the Landau pole contribution becomes more dominant. In this region the effect of the non-universalities tends to get washed out. Further the effect of the Z-pole disappears for $m_{\tilde{\chi}_1} \geq 65$ GeV. This explains in part why the minimum and the maximum event rates in the region of large gluino masses are insensitive to the effect of the non-universalities. Also, as m_t is increased, D_0 becomes smaller and the Landau pole term in Eq. (40) dominates, making the effects of non-universal terms again less significant. Numerical analysis supports this conclusion.

Finally we discuss the consequences of a more accurate determination of Ωh^2 . One expects such a situation to occur in the next round of COBE like sat-




Figure 6: Maximum and minimum of event rates for Xenon for the non-universal case when $\delta_1 = -1 = -\delta_2$

telite experiments which may determine the Hubble parameter to within 10% or better accuracy. This determination of the Hubble parameter will strongly reduce the error in Ωh^2 . Here we analyse what such an accurate determination will do to the prediction of the neutralino event rates. For specificity let us assume that Ωh^2 is constrained to lie in the narrow band $0.225 < \Omega h^2 < 0.275$. The narrowing of the range of Ωh^2 more stringently limits the parameter space of the SUGRA model. The result of the event rate analysis for Xenon under the above constraint is shown in Fig.9. An interesting new phenomenon is the appearance of peaks in the minimum curves. These sharp peaks originate from the constraints to prevent charge and color breaking minima⁵⁰ (CCB constraints). The position of the peak is determined by the mass parameter $m_{H_2}^2$. The peak moves to the right as $m_{H_2}^2$ increases and moves to the left as $m_{H_2}^2$ decreases. The first situation occurs for the case $\delta_1 < 0, \delta_2 > 0$, and the second for the case $\delta_1 > 0, \delta_2 < 0$. Numerical analysis for these cases as given by the dashed curve and the dotted curve support the analytical result. Comparison of Figs.8 and 9 shows that a narrower range in Ωh^2 more sharply constrains the SUGRA parameter space and the allowed band of event rates shrinks.

9 CONCLUDING REMARKS

In this review we have given an analysis of event rates for the observation of dark matter in terrestrial dark matter detectors. Supergravity GUT models under the assumption of R parity invariance predict the neutralino to be cold dark matter over most of the parameter space of the model. The analysis of the event rates is carried out by ranging over the full parameter space of the SUGRA model within the prescribed naturalness constraints. Then we find that the event rates can range from $R = (10^{-5} \text{ to } \approx 5) \text{ events/kg da.}$




Figure 7: Maximum and minimum of event rates for Xenon for the non-universal case when $\delta_1 = 1 = -\delta_2$

We find that for a heavy target such as Ge or Xenon the maximum event rates are almost totally coherent (which is controlled by the scalar interaction) while the minimum event rates can have a significant incoherent component (which is controlled by the spin dependent interaction). For a light target such as CaF_2 , one finds that the minimum event rates are essentially incoherent while the maximum event rates have a very significant component of coherent scattering.

We have also analysed in this paper the effects of non-universal soft SUSY breaking parameters in the Higgs sector on event rates. It is found that the non-universal soft breaking effects are most significant in the neutralino mass range $m_{\tilde{\chi}_1} \simeq (35\text{--}65)$ GeV. In this region one finds that the minimum event rates can be decreased or increased by a factor of 10 depending on the nature of non-universality.

The above analyses show that experiments with a sensitivity of $R \geq 0.005$ events/kg da can explore a significant part of the parameter space. However, a further significant improvement in sensitivity by a factor of about 100 or more would be needed to explore the entire parameter space. For the case of non-universality with $\delta_2 = -\delta_1 = 1$ there is an improvement by a factor of about 10 in the event rates in the energy domain $M_{\tilde{Z}_1} \leq 65$ GeV. However, for the case $\delta_2 = -\delta_1 = -1$ one finds a reduction in the event rates by a factor of about 10 in the same mass region. Thus whether or not the event rates are enhanced or reduced depends on the nature of the non-universality.

The above results have important implications for dark matter detectors. Thus the analysis shows that the parameter space of the model with $\delta_2 = -\delta_1 = 1$ can be exhausted completely in the mass domain $M_{\tilde{Z}_1} \leq 65$ GeV by an improvement in the detector sensitivity by a factor of about 100.




Figure 8: Comparison of the maximum and minimum of event rates for Xenon for three cases:(i) universal case when $\delta_1 = 0 = \delta_2$ (solid); (ii)(i)non-universal case when $\delta_1 = -1 = -\delta_2$ (dashed); (iii) non-universal case when $\delta_1 = 1 = -\delta_2$ (dotted)

However, much greater sensitivity will be needed to do the same for the case $\delta_2 = -\delta_1 = 1$. Thus new detection techniques which have the potential of dramatically improving the detection sensitivities^{27,51}, are important for a complete exploration of the parameter space of the supergravity unified model.

Acknowledgments

This research was supported in part by NSF grant numbers PHY-96020274 and PHY-9411543.

Appendix A

To second order perturbation theory one finds³³ when $(M_Z/\mu)^2 \ll 1$:

$$n_1 \cong -\frac{1}{2} \frac{M_Z}{\mu} \frac{1}{(1 - \tilde{m}_1^2/\mu^2)} \frac{M_Z}{\tilde{m}_2 - \tilde{m}_1} \sin 2\theta_W \left[\sin 2\beta + \frac{\tilde{m}_1}{\mu} \right] \quad (48)$$

$$n_2 = 1 - \frac{1}{2} \frac{M_Z^2}{\mu^2} \frac{1}{(1 - \tilde{m}_1^2/\mu^2)^2} \sin^2 \theta_W \left[1 + \frac{\tilde{m}_1}{\mu} \sin 2\beta + \frac{\tilde{m}_1^2}{\mu^2} \right] \quad (49)$$




Figure 9: Comparison of the maximum and minimum of event rates for Xenon for three cases when $0.225 < \Omega h^2 < 0.275$: (i) universal case when $\delta_1 = 0 = \delta_2$ (solid); (ii) non-universal case when $\delta_1 = -1 = -\delta_2$ (dashed); (iii) non-universal case when $\delta_1 = 1 = -\delta_2$ (dotted)

$$n_3 = \frac{M_Z}{\mu} \frac{1}{1 - \tilde{m}_1^2/\mu^2} \sin\theta_W \sin\beta \left[1 + \frac{\tilde{m}_1}{\mu} \cot\beta \right] \quad (50)$$

$$n_4 = -\frac{M_Z}{\mu} \frac{1}{1 - \tilde{m}_1^2/\mu^2} \sin\theta_W \cos\beta \left[1 + \frac{\tilde{m}_1}{\mu} \tan\beta \right] \quad (51)$$

Eqs. (48-51) differ from the numerical computer results by amounts $\delta n_i \lesssim 0.03$ over a large part of the parameter space and are often significantly better.

Appendix B

As discussed in Sec 3, the conventional approximation in the computation of relic density has been the expansion of σv in a power series expansion in v . However, in the annihilation of the neutralinos one encounters Z and Higgs poles. A power series expansion of σv in this case breaks down. One must therefore treat the pole correctly in the thermal averaging and in the computation of $J(x_f)^{15}$. We illustrate this below for the case of the Higgs pole. Here one has

$$\sigma v = \frac{A_{Higgs}}{m_{\chi_1}^4} \frac{v^2}{((v^2 - \epsilon_h)^2 + \gamma_h^2)} \quad (52)$$

$$\epsilon_h = (m_h^2 - 4m_{\chi_1}^2)/m_{\chi_1}^2 \quad (53)$$

$$\gamma_h = m_h \Gamma_h / m_{\chi_1}^2 \quad (54)$$

In the above m_h is the Higgs mass and Γ_h is the Higgs width. $J(x_f)$ involves a double integration over a pole which for accuracy requires a high precision numerical computation. It is possible to significantly simplify the expression for $J(x_f)$ by carrying out one of the integrations analytically. One may then write $J(x_f)$ in the following form

$$J_{Higgs}(x_f) = \frac{A_{Higgs}}{2\sqrt{2}m_{\chi_1}^4} (I_{1h} + \frac{\epsilon_h}{\gamma_h} I_{2h}) \quad (55)$$

$$I_{1h} = \frac{1}{2} \int_0^\infty dy y^{-\frac{1}{2}} e^{-y} \log \left[\frac{(4yx_f - \epsilon_h)^2 + \gamma_h^2}{\epsilon_h^2 + \gamma_h^2} \right] \quad (56)$$

$$I_{2h} = \frac{1}{2} \int_0^\infty dy y^{-\frac{1}{2}} e^{-y} \left[\tan^{-1} \left(\frac{(4yx_f - \epsilon_h)^2 + \gamma_h^2}{\gamma_h} \right) + \tan^{-1} \left(\frac{\epsilon_h}{\gamma_h} \right) \right] \quad (57)$$

The numerical analysis of $J(x_f)$ is significantly simplified using the reduced form. A similar treatment of the Z pole term can also be carried out.

References

1. E.W. Kolb and M.S. Turner, "The Early Universe" (Addison-Wesley, Redwood City, 1989).
2. P.F. Smith and J.D. Lewin, Phys.Rep.187,No.5,203(1990).
3. G. Jungman, M. Kamionkowski and K. Greist, "Supersymmetric Dark Matter", Phys. Rep. **267**,195(1995).
4. J. Primack et al. Phys. Rev. Lett. **74**, 2160 (1995).
5. A.H. Chamseddine, R. Arnowitt and P. Nath, Phys. Rev. Lett. **29**, 970 (1982). For reviews see P.Nath, R. Arnowitt and A.H.Chamseddine, "Applied N = 1 Supergravity" (World Scientific, Singapore, 1984); H.P. Nilles, Phys. Rep. **110**, 1 (1984); R. Arnowitt and P. Nath, Proc. of VII J.A. Swieca Summer School ed. E. Eboli (World Scientific, Singapore, 1994).
6. L.Hall, J.Lykken and S.Weinberg, Phys. Rev. **D27**,2359(1983); P.Nath, R.Arnowitt and A.H.Chamseddine, Nucl.Phys. **B227**,121(1983).
7. K. Inoue et al. Prog. Theor. Phys **68**, 927 (1982); L. Ibañez and G.G. Ross, Phys. Lett. **B110**, 227 (1982); L. Avarez-Gaumé, J.Polchinski and M.B. Wise, Nucl. Phys. **B221**, 495 (1983), J. Ellis, J. Hagelin, D.V.

- Nanopoulos and K. Tamvakis, Phys. Lett. **B125**, 2275 (1983); L.E. Ibañez and C. Lopez, Nucl. Phys. **B233**, 545 (1984); L.E. Ibañez, C. Lopez and C. Muñoz, Nucl. Phys. **B256**, 218 (1985).
8. J. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. **B262**, 477 (1991); R.Arnowitt and P.Nath, Phys.Rev.**D46**,3981(1992).
 9. For more recent analyses of supergravity unification with radiative breaking of the electro-weak symmetry see, G.Ross and R. Roberts, Nucl.Phys.**B377**, 571(1992); R.Arnowitt and P.Nath, Phys.Rev. Lett. **69**,725(1992); P.Nath and R.Arnowitt, Phys.Lett.**B289**,368(1992); M.Drees and M.M. Nojiri, Nucl.Phys.**B369**,54(1993); S.Kelley, J.Lopez, D.Nanopoulos, H.Pois and K. Yuan, Nucl.Phys.**B398**,3(1993); M. Olechowski and S. Pokorski, Nucl. Phys.**B404**,590(1993); V.Barger, M.Berger, and P. Ohman, Nucl.Phys. **B49**, 4908(1994); G. Kane, C. Kolda, L. Roszkowski and J. Wells, Phys. Rev. **D49**, 6173 (1994); D.J. Castano, E. Piard, and P.Ramond, Phys.Rev.**D49**,4882(1994); W.de Boer, R. Ehret, and D. Kazakov, Z. Phys.**C67**,647(1995); H.Baer, C.H.Chen, R. Munroe, F.Paige, F.Paige, and X.Tata, Phys.Rev.**D51**,1046(1995).
 10. L.Ibanez, C.Lopez, and C.Munos, Nucl. Phys.**B256**, 218(1985).
 11. P. Nath, J. Wu and R. Arnowitt, Phys.Rev.**D52**,4169(1995).
 12. B.W. Lee and S. Weinberg, Phys. Rev. Lett. **39**, 165 (1977); D.A. Dicus, E. Kolb and V. Teplitz, Phys. Rev. Lett. **50**, 168 (1977); H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983); J. Ellis, J.S. Hagelin, D.V. Nanopoulos and N. Srednicki, Nucl. Phys. **B238**, 453 (1984); J.Lopez, D.V.Nanopoulos and K. Yuan, Nucl.Phys.**B370**,445(1992); M. Drees and M.M. Nojiri, Phys. Rev. **D47**,376(1993).
 13. K. Greist and D. Seckel, Phys. Rev. **D43**, 3191 (1991); P. Gondolo and G. Gelmini, Nucl. Phys. **B360**, 145 (1991).
 14. R. Arnowitt and P. Nath, Phys. Lett. **B299**, 58 (1993); **B303**, 403 (1993) (E).
 15. P. Nath and R. Arnowitt, Phys. Rev. Lett. **70**, 3696 (1993).
 16. H.Baer and M.Brhllick, Phys.Rev.**D53**,597(1996).
 17. M.S. Alam et al. (CLEO Collaboration), Phys. Rev. Lett. **74**, 2885 (1995).
 18. S. Bertolini, F. Borzumati and A. Masiero, Phys. Rev. Lett. **59**, 180 (1987); B. Grinstein, R. Springer and M.B. Wise, Nucl. Phys. **B339**, 269 (1990); S. Bertolini, F. Borzumati, A. Masiero and G. Ridolfi, Nucl. Phys. **B353**, 591 (1991); R. Barbieri and G. Giudice, Phys. Lett. **B309**, 86 (1993); M. Misiak, Phys. Lett. **B269**, 161 (1991); Nucl. Phys. **B393**, 23 (1993).

19. A.J. Buras, M. Misiak, M. Münz and S. Pokorski, Nucl. Phys. **B424**, 374 (1994); M. Ciuchini, E. Franco, G. Martinelli, L. Reina and L. Silvestrini, Phys. Lett. **B316**, 127 (1993).
20. P.Nath and R.Arnouitt, Phys.Lett.**B336**, 395(1994).
21. F.Borzumati, M.Drees and M.Nojiri, Phys.Rev. **D51**,341(1995).
22. M. Misiak, talk at ICHEP96, Warsaw, Poland, July 1996.
23. H.Baer and M.Brhllick, hep-ph/9610224.
24. M.Kamionkowski, Phys.Rev.**D44**,3021(1991); F.Halzen, T.Saltzer and M. Kamionkowski, Phys.Rev. **D45**,139(1992);F.Halzen, talk at this workshop.
25. M. Kamionkowski, K.Greist, G.Jungman, and B.Sadoulet, Phys.Rev.Lett. **74**,5174(1995).
26. G.D. Starkman and D.N. Spergel, Phys.Lett.**74**,2623(1995).
27. J.I. Collar, Phys.Rev.**D54**,1247(1996); L.Lessard, talk at this conference.
28. M.W. Goodman and E. Witten, Phys. Rev. **D31**, 3059(1983); K.Greist, Phys. Rev. **D38**, (1988)2357; **D39**,3802(1989)(E); J.Ellis and R.Flores, Phys. Lett. **B300**,175(1993); R. Barbieri, M. Frigeni and G.F. Giudice, Nucl. Phys. **B313**,725(1989); M.Srednicki and R.Watkins, Phys. Lett. **B225**,140(1989); R.Flores, K.Olive and M.Srednicki, Phys. Lett. **B237**,72(1990).
29. A. Bottino etal, Astro. Part. Phys. **1**, 61 (1992); **2**, 77 (1994).
30. V.A. Bednyakov, H. V. Klapdor-Kleingrothaus and S.G. Kovalenko, Phys. Rev. **D50**,7128(1994).
31. R.Arnouitt and P.Nath, Mod. Phys. Lett. **A 10**,1257(1995).
32. P.Nath and R.Arnouitt, Phys. Rev. Lett.**74**,4592(1995).
33. R.Arnouitt and P.Nath, Phys. Rev. **D54**,2374(1996).
34. E.Diehl, G.Kane, C.Kolda and J.Wells, Phys. Rev.**D52**,4223(1995).
35. L. Bergstrom and P.Gondolo, hep-ph/9510252.
36. J.D. Vergados, J. Phys.**G22**,253(1996).
37. G.L.Kane and J.D. Wells, Phys.Rev. Lett.**76**, 4458(1996).
38. B. de Carlos and G.V. Kraniotis, hep-ph/9610355.
39. J.F. Gunion, H.E. Haber, G. Kane and S. Dawson, "The Higgs Hunter's Guide" (Addison-Wesley Publ Co., Redwood City, 1990).
40. H. Haber and R. Hempfling, Phys. Rev. Lett. **60**, 1815 (1991); Phys. Rev. **D48**, 4280 (1993).
41. M.Kamionkowski, Phys.Rev.**D44**,3021(1991).
42. T.P. Cheng, Phys. Rev. **D38**, 2869 (1989); J. Gasser, H. Leutwyler and M.E. Saino, Phys. Lett. **B253**, 252 (1991).
43. J. Ashman etal (EMC Collaboration), Nucl. Phys. **B328**, 1 (1989); A. Manohar and R. Jaffe, Nucl. Phys. **B337**, 509 (1990).

44. D. Adams et al. (SMC Collaboration), Phys. Lett. **B329**, 399 (1994);
R. Arnold et al. (E143 Collaboration) in “Intersections of Particle and
Nuclear Physics”, St. Petersburg, FL (1994); J.Ellis and M.Karliner,
Phys.Lett.**B341**,397(1995).
45. S.K.Soni and H.A. Weldon, Phys.Lett.**B126**,215(1983);V.S.Kaplunovsky
and J.Louis, Phys.Lett.**B306**, 268(1993).
46. D.Matalliotakis and H.P.Nilles, Nucl.Phys.**B435**, 115(1995).
47. M.Olechowski and S.Pokorski, Phys.Lett. **B344**, 201(1995).
48. Polonski and Pomerol,Phys.Rev.**D51**,6532(1995).
49. V.Berezinsky, A.Bottino, J.Ellis, N.Forrengo, G.Mignola, and S.Scopel,
Astropart.Phys.5:1-26,1996.
50. J.M. Frere, D.R.T. Jones and S.Raby, Nucl. Phys.**B222**, 11(1983);
M.Drees, M.Gluck and K.Grassie, Phys. Lett.**B157**, 164(1985); J.A.
Casas, A. Lleyda and C. Munoz, Nucl. Phys. **B471**, 3(1996); see
however, A. Kusenko, P. Langacker, and G. Segre, Phys. Rev.**D54**,
5824(1996).
51. D.Cline.”On a Discriminatory Liquid Xenon SUSY Wimp Detector”,
Nucl. Phys. B (in press); P.Benetti et al, Nucl. Inst. and Method
for Particle Physics Research, **A307** (1993) 203; See also talks by R.
Bernabei, D.Cline, and H.Wang at this conference.